



**The Cochrane Library.** The single most reliable source of evidence in healthcare

The Cochrane Reviews highlighted below are available from the

Cochrane Database of Systematic Reviews ([www.thecochranelibrary.com](http://www.thecochranelibrary.com))

Cochrane Reviews are regularly updated as new evidence emerges and in response to feedback, and the Cochrane Database of Systematic Reviews should always be consulted for the most recent version of the review.

## STUDY REPORTS PATIENT-REPORTED LOSS OF SMELL IN 86 PERCENT OF MILD COVID-19 CASES

**06 JANUARY 2021 - WILEY** - A reduced sense of smell, or olfactory dysfunction, is one of the most common symptoms of COVID-19. A recent study published in the *Journal of Internal Medicine* has examined its prevalence and recovery in patients with varying degrees of severity of COVID-19.

In the study of 2,581 patients from 18 European hospitals, the patient-reported prevalence of olfactory dysfunction was 85.9% in mild cases of COVID-19, 4.5% in moderate cases, and 6.9% in severe-to-critical cases. The average duration of olfactory dysfunction reported by patients was 21.6 days, but nearly one-quarter of affected patients reported that they did not recover their sense of smell 60 days after losing it.

Objective clinical evaluations identified olfactory dysfunction in 54.7% of mild cases of COVID-19 and 36.6% of moderate-to-critical cases of COVID-19. At 60 days and 6 months, 15.3% and 4.7% of these patients did not objectively recover their sense of smell, respectively.

“Olfactory dysfunction is more prevalent in mild COVID-19 forms than in moderate-to-critical forms, and 95% of patients recover their sense of smell at 6-months post-infection,” said lead author Jerome R. Lechien, MD, PhD, MS, of Paris Saclay University.

### Additional Information

**Link to Study:** <https://onlinelibrary.wiley.com/doi/10.1111/joim.13209>

*Copyright © 2021 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd., reproduced with permission.*